

Seesaw parametrization for n right-handed neutrinos

Julian Heeck*

Max-Planck-Institut für Kernphysik, Saupfercheckweg 1, 69117 Heidelberg, Germany

Introducing n right-handed neutrinos to the Standard Model yields, in general, massive active neutrinos. We give explicit parametrizations for the involved mixing and coupling matrices in terms of physical parameters for both the top-down and the bottom-up approach, for arbitrary n . Bounds on the complex mixing angles in the bottom-up approach from perturbativity of the Yukawa couplings to charged lepton flavor violation are discussed. As a novel illustration of possible effects from $n \neq 3$, we extend the neutrino anarchy framework to arbitrary n ; we show that while the anarchic mixing angles are insensitive to the number of singlets, the observed ratios of neutrino masses prefer small n for the simplest linear measure.

PACS numbers: 14.60.Pq, 14.60.St

Keywords: Neutrino Physics, Beyond Standard Model

I. INTRODUCTION

The (type-I) seesaw mechanism [1] is arguably the simplest and best motivated framework to not only give neutrinos mass, but also explain the smallness of said masses compared to the electroweak scale. The necessary right-handed neutrino partners are assumed to be heavy to suppress the active neutrino masses, which also allows for baryogenesis through leptogenesis [2] via the decay of the heavy states, thus explaining the baryon asymmetry of the universe. While initially motivated in the context of grand unified theories, the seesaw mechanism has since been applied in its own right. The number of right-handed neutrinos, transforming as singlets of the Standard Model (SM) gauge group, is then no longer constrained by anomaly cancellations or the need to fill up an irreducible representation, but rather a free parameter. At least two right-handed neutrinos are necessary to reproduce the observed neutrino mass-squared differences, as well as leptogenesis [3]. There is however no upper bound, and models with $\mathcal{O}(10^2\text{--}10^3)$ singlets have been studied in the context of leptogenesis [4], lepton flavor violation (LFV) [5] and as a way to explain the large mixing angles of the active neutrinos [6]. Like for most things in life, there is also a motivation by string theory [7]. The number of singlets is formally infinite in extra-dimensional theories, as right-handed neutrinos are not restricted to the SM-brane and will thus lead to a tower of Kaluza-Klein excitations [8].

Still lacking, however, is a proper parametrization of the arising mixing and coupling matrices in terms of physical quantities, as needed for example for efficient parameter scans (as the number of redundant parameters explodes for large n). The goal of this work is to provide exactly these parametrizations for arbitrary n , both for the most general case (top-down approach, to some degree discussed in Refs. [9]) and the seesaw limit (bottom-up).

The rest of this paper is organized as follows: We first fix our notation in Sec. II, discuss the top-down parametrization in Sec. III and the bottom-up parametrization in Sec. IV, also deriving constraints on the involved parameters from perturbativity and LFV. To illustrate possible effects of n singlets we discuss basis independence (neutrino anarchy) in this framework and present the resulting distributions for the neutrino masses in Sec. V. App. A gives a brief introduction to the group $U(N)$ and various representations of its elements, as needed for our parametrizations. App. B collects and extends basis independent measures for miscellaneous matrices necessary for the discussion of anarchy. Finally, App. C is devoted to the somewhat special case $n = 2$, which would interrupt the flow if included in the main text, which will only cover $n \geq 3$.

*Electronic address: julian.heeck@mpi-hd.mpg.de

II. FRAMEWORK AND NOTATION

Introducing n SM singlet right-handed neutrinos N_j to the Standard Model modifies the Lagrangian by the following terms:

$$\mathcal{L} = \mathcal{L}_{\text{SM}} + i\overline{N}_j \not{\partial} N_j - \left(\overline{N}_j (\mathbf{Y}_\nu)_{j\alpha} \tilde{H}^\dagger L_\alpha + \frac{1}{2} \overline{N}_j (\mathcal{M}_R)_{jk} N_k^c + \text{h.c.} \right), \quad (1)$$

where a sum over $\alpha = e, \mu, \tau$ (sometimes 1, 2, 3 in the following) and $j, k = 1, 2, \dots, n$ is understood and will in the following often be denoted in a vector notation (e.g. as $\overline{N}\mathcal{M}_R N^c$). Without loss of generality we work in a basis where the charged lepton mass matrix is diagonal, which can be accomplished by unitary transformations of the lepton fields. \mathcal{M}_R can be diagonalized by an $n \times n$ unitary matrix V like $V^T \mathcal{M}_R V = \text{diag}$. This merely redefines \mathbf{Y}_ν , so we will from now on work in a basis where \mathcal{M}_R is diagonal, with positive real entries M_i .[‡] After the Higgs doublet H acquires its vacuum expectation value v , the Dirac mass matrix for the neutrinos $m_D = v\mathbf{Y}_\nu$ is generated, leading to a $(3+n) \times (3+n)$ Majorana mass matrix for the neutral fermions $(\nu_L, N^c)^T$:

$$\mathcal{M}_{\text{full}} = \begin{pmatrix} 0 & m_D^T \\ m_D & \mathcal{M}_R \end{pmatrix}. \quad (2)$$

Assuming $\mathcal{M}_R \gg m_D$ gives the low-energy neutrino mass matrix for the flavor neutrinos $\nu_f \simeq \nu_L - m_D^\dagger \mathcal{M}_R^{-1} N^c$ of seesaw form:

$$\mathcal{M}_\nu \simeq -m_D^T \mathcal{M}_R^{-1} m_D = -v^2 \mathbf{Y}_\nu^T \mathcal{M}_R^{-1} \mathbf{Y}_\nu. \quad (3)$$

Diagonalization of \mathcal{M}_ν can be performed in the following way

$$\mathcal{M}_\nu = U_{\text{PMNS}}^* \text{diag}(m_1, m_2, m_3) U_{\text{PMNS}}^\dagger \equiv U_{\text{PMNS}}^* d_m U_{\text{PMNS}}^\dagger, \quad (4)$$

with the unitary Pontecorvo-Maki-Nakagawa-Sakata matrix (PMNS matrix) in standard parametrization

$$U_{\text{PMNS}} = P' \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta} \\ -c_{23}s_{12} - s_{23}s_{13}c_{12}e^{i\delta} & c_{23}c_{12} - s_{23}s_{13}s_{12}e^{i\delta} & s_{23}c_{13} \\ s_{23}s_{12} - c_{23}s_{13}c_{12}e^{i\delta} & -s_{23}c_{12} - c_{23}s_{13}s_{12}e^{i\delta} & c_{23}c_{13} \end{pmatrix} P, \quad (5)$$

and the Majorana phase matrix $P = \text{diag}(1, e^{i\alpha/2}, e^{i\beta/2})$. Here we used the abbreviations $s_{ij} \equiv \sin \theta_{ij}$ and $c_{ij} \equiv \cos \theta_{ij}$ for the three mixing angles. The phase matrix $P' = \text{diag}(e^{ia/2}, e^{ib/2}, e^{ic/2})$ can be absorbed by the lepton fields and is hence unphysical. The neutrino mass eigenstates are then $\nu_m = U_{\text{PMNS}}^\dagger \nu_f$.

At low energies the 9 complex entries of the symmetric Majorana matrix \mathcal{M}_ν decompose into the three eigenvalues m_i (neutrinos masses), three mixing angles $\theta_{23}, \theta_{12}, \theta_{13}$ (atmospheric, solar and reactor angle) and three CP violating phases δ, α, β , the latter two being unobservable in neutrino oscillations (but in principle testable in $0\nu\beta\beta$ experiments [10]).

III. TOP-DOWN PARAMETRIZATION

All the information about neutrino mixing is encoded in \mathbf{Y}_ν , the only non-diagonal matrix in the lepton sector. For the general complex $n \times 3$ matrix \mathbf{Y}_ν , we can write down a singular value decomposition

$$\mathbf{Y}_\nu = V_R \begin{pmatrix} \text{diag}(y_1, y_2, y_3) \\ \vec{0}_3 \\ \vdots \\ \vec{0}_3 \end{pmatrix} V_L^\dagger \equiv V_R D_Y V_L^\dagger, \quad (6)$$

[‡] This is not only convenient, but also necessary to eliminate unphysical parameters. To this effect we note that a diagonal \mathcal{M}_R can store $\mathcal{O}(n)$ parameters (together with \mathbf{Y}_ν), while a non-diagonal \mathcal{M}_R has $\mathcal{O}(n^2)$ independent entries.

with positive singular values y_i . Out of the $6n$ real parameters in \mathbf{Y}_ν , 3 phases can be absorbed in the lepton fields, so only $6n - 3$ are physical. We will confirm this with an explicit parametrization of the unitary matrices $V_{R,L}$ below. For the 3×3 matrix V_L , we take the PMNS parametrization from Eq. (5), while the $n \times n$ matrix V_R can be written as

$$V_R = \left(\prod_{i=1}^n \prod_{j=i+1}^n \Omega_{ij}(\alpha_{ij}, \xi_{ij}) \right) \times \text{diag}(e^{i\phi_1}, \dots, e^{i\phi_n}). \quad (7)$$

Here, $\Omega_{ij}(\alpha_{ij}, \xi_{ij})$ denotes matrices of the form

$$\Omega_{ij}(\alpha_{ij}, \xi_{ij}) \equiv \begin{pmatrix} 1 & & & & \\ & \ddots & & & \\ & & \cos(\alpha_{ij}) & \sin(\alpha_{ij})e^{i\xi_{ij}} & \\ & & -\sin(\alpha_{ij})e^{-i\xi_{ij}} & \cos(\alpha_{ij}) & \\ & & & & \ddots & \\ & & & & & 1 \end{pmatrix}, \quad (8)$$

where $\sin(\alpha_{ij})e^{i\xi_{ij}}$ sits at the i -th row and j -th column. See App. A for a derivation and a proof for the validity of this form of V_R . It is easy to check that V_R (V_L) has the proper n^2 (3^2) real parameters, $n(n+1)/2$ (6) of which are phases. From the form of \mathbf{Y}_ν in Eq. (6), it is clear that the angles ϕ_j either act on zeros in D_Y or can be absorbed by the phase matrix P_L^\dagger in V_L^\dagger . An overall phase and $P_L'^\dagger$ can be absorbed by the lepton fields, so V_L contains only 3 angles and 3 phases. The ordering of the Ω_{ij} in V_R is of course arbitrary, which allows us to move all the rotations that act on the zeros in D_Y to the right, so $(n-3)(n-4)/2$ matrices of the type (8) drop out, leaving $n(n-1) - (n-3)(n-4) = 6n - 12$ real parameters in V_R , half of which are phases. Consequently we can restrict the product in Eq. (7) to $\prod_{i=1}^3 \prod_{j=i+1}^n$.[§] As expected, \mathbf{Y}_ν contains 3 (from y_i) plus 6 (V_L) plus $6n - 12$ (V_R) equals $6n - 3$ real, physical parameters, $3(n-1)$ of which are phases. This agrees with the parameter counting in Ref. [11].

The full mass matrix can then be written as

$$\mathcal{M}_{\text{full}} = \begin{pmatrix} 0 & vV_L^* D_Y^T V_R^T \\ vV_R D_Y V_L^\dagger & \mathcal{M}_R \end{pmatrix}, \quad (9)$$

with the explicit form for V_L :

$$V_L = \begin{pmatrix} c_{12}^L c_{13}^L & s_{12}^L c_{13}^L & s_{13}^L e^{-i\beta_1^L} \\ -c_{23}^L s_{12}^L - s_{23}^L s_{13}^L c_{12}^L e^{i\beta_1^L} & c_{23}^L c_{12}^L - s_{23}^L s_{13}^L s_{12}^L e^{i\beta_1^L} & s_{23}^L c_{13}^L \\ s_{23}^L s_{12}^L - c_{23}^L s_{13}^L c_{12}^L e^{i\beta_1^L} & -s_{23}^L c_{12}^L - c_{23}^L s_{13}^L s_{12}^L e^{i\beta_1^L} & c_{23}^L c_{13}^L \end{pmatrix} \text{diag}(1, e^{i\beta_2^L/2}, e^{i\beta_3^L/2}), \quad (10)$$

and the shorthand notation $s_{ij}^L = \sin \theta_{ij}^L$ etc. for the mixing angles. In the seesaw limit, we thus find

$$\mathcal{M}_\nu \simeq -v^2 V_L^* D_Y^T (V_R^\dagger \mathcal{M}_R V_R^*)^{-1} D_Y V_L^\dagger. \quad (11)$$

Note that the occurring mass matrix $V_R^\dagger \mathcal{M}_R V_R^*$ is for $n > 3$ no longer the most general complex symmetric $n \times n$ matrix, because we already restricted V_R to the physical angles.

From Eq. (9) or Eq. (11) one can now extract the physical masses and mixing angles in terms of the parameters y_i , α_{ij} , ξ_{ij} , θ_{ij}^L and β_i^L .

For completeness, let us count the total number of parameters in the lepton sector. We have 3 charged lepton masses, n heavy neutrino masses and $6n - 3$ real parameters in \mathbf{Y}_ν , summing up to $7n$ real parameters.

[§] In other words, the $U(n-3)$ subgroup of $U(n)$, acting on the lower right parts, can be modded out.

IV. BOTTOM-UP PARAMETRIZATION

Seeing as the seesaw formula (11) contains more parameters than measurable observables, it proves convenient to parameterize our ignorance of the high-energy sector of the seesaw model. In other words, there are infinitely many different \mathbf{Y}_ν and \mathcal{M}_R that lead to the same low-energy neutrino parameters via the seesaw formula $\mathcal{M}_\nu \simeq -m_D^T \mathcal{M}_R^{-1} m_D$. The parametrization of this ambiguity is the goal of the Casas-Ibarra-parametrization [12] of the Yukawa coupling:

$$\mathbf{Y}_\nu = \frac{i}{v} \sqrt{\mathcal{M}_R} R \sqrt{d_m} U_{\text{PMNS}}^\dagger. \quad (12)$$

This \mathbf{Y}_ν solves

$$-v^2 \mathbf{Y}_\nu^T \mathcal{M}_R^{-1} \mathbf{Y}_\nu \simeq \mathcal{M}_\nu = U_{\text{PMNS}}^* d_m U_{\text{PMNS}}^\dagger, \quad (13)$$

as long as the complex $n \times 3$ matrix R satisfies $R^T R = \mathbb{1}_{3 \times 3}$. Correspondingly, any values of \mathcal{M}_R and R in Eq. (12) lead to the same low-energy model, which means we can fix the PMNS mixing angles and masses to the observed ones and study \mathbf{Y}_ν independently in other processes.

We will now give an explicit parametrization of R , which has $3n$ complex entries obeying 12 real constraints ($R^T R = \mathbb{1}$ is symmetric); we therefore expect to find $6(n-2)$ real physical parameters. This agrees of course with the parameter counting in the last section, as we also have 3 charged lepton masses, 3 light neutrino masses, n heavy neutrino masses and 6 parameters in the PMNS matrix, summing up to $7n$ input parameters. Since we only need three orthonormal complex vectors $\in \mathbb{C}^n$ to define R , we can write

$$R = O_{n \times n} \begin{pmatrix} \mathbb{1}_{3 \times 3} \\ \vec{0}_3 \\ \vdots \\ \vec{0}_3 \end{pmatrix} \equiv O_{n \times n} \mathbb{1}_{n \times 3}, \quad (14)$$

with a complex orthogonal $n \times n$ matrix $O_{n \times n} \in O(n, \mathbb{C})$. Similar to the $U(n)$ parametrization used for V_R in the previous section, we can write $O_{n \times n}$ as a product of the $\Omega_{ij}(\gamma_{ij}, \xi'_{ij})$ matrices from Eq. (8), but with the arguments $\xi'_{ij} = 0$ and $\gamma_{ij} \in \mathbb{C}$ in order for $O_{n \times n}$ to be complex orthogonal:

$$O_{n \times n} = S \times \prod_{i=1}^n \prod_{j=i+1}^n \Omega_{ij}(\gamma_{ij}, 0). \quad (15)$$

Here, S is a diagonal matrix with entries ± 1 , which decides the determinant of $O_{n \times n}$ and therefore whether it describes a rotation or a reflection. We can once again mod out the $O(n-3, \mathbb{C})$ subgroup that acts on the zeros in $\mathbb{1}_{n \times 3}$ by reordering the rotations, so the product in Eq. (15) can be restricted to $\prod_{i=1}^3 \prod_{j=i+1}^n$. Generally speaking, R is an element of the quotient space $O(n, \mathbb{C})/O(n-3, \mathbb{C})$. For real angles γ_{ij} , R is an element of the compact Stiefel manifold $V_3(\mathbb{R}^n) \cong O(n, \mathbb{R})/O(n-3, \mathbb{R})$, which admits a Haar measure and could thus be used for statistical considerations.

Note that the case $n = 4$ is special in the sense that there are no superfluous rotations that need to be modded out. One can therefore take any complete parametrization for $O_{4 \times 4}$ without having to worry about unphysical parameters.

For complex angles, $O_{n \times n}$ and therefore R and \mathbf{Y}_ν can have arbitrarily large elements, because the entries $\cos ix = \cosh x$ grow exponentially with x . As a result, the Yukawa couplings \mathbf{Y}_ν can be very large even for small \mathcal{M}_R , which is somewhat counterintuitive to the seesaw formula and relies on cancellations in the matrix product $\mathbf{Y}_\nu^T \mathcal{M}_R^{-1} \mathbf{Y}_\nu$. Even though these cancellations suggest finetuning of parameters, this is a viable possibility and deserves discussion, especially because a low \mathcal{M}_R with large Yukawa couplings makes an LHC discovery (direct or indirect) possible [13]. Popular models to obtain this finetuning in a natural way are for example inverse- [14] and linear seesaw [15], usually based on the case $n = 6$ with an imposed structure in \mathcal{M}_R and \mathbf{Y}_ν by some symmetry.

A useful constraint on the complex mixing angles γ_{ij} comes from the perturbativity of \mathbf{Y}_ν , on which many calculations involving \mathbf{Y}_ν rely on. For this, we propose yet another, slightly different parametrization of R , which we obtain by noting that $\Omega_{ij}(\gamma_{ij}, 0) = \Omega_{ij}(\text{Re}(\gamma_{ij}), 0) \times \Omega_{ij}(i\text{Im}(\gamma_{ij}), 0)$, and that the

ordering of the Ω_{ij} is arbitrary:

$$R = S \times \left(\prod_{i=1}^3 \prod_{j=i+1}^n \Omega_{ij}(\eta_{ij}, 0) \right) \left(\prod_{i=1}^3 \prod_{j=i+1}^n \Omega_{ij}(i\rho_{ij}, 0) \right) \mathbb{1}_{n \times 3}. \quad (16)$$

The $6(n-2)$ parameters that make up R are thus equally divided among the new real parameters η_{ij} and ρ_{ij} . The product involving η_{ij} is just an orthogonal matrix, so we can restrict $\eta_{ij} \in [0, 2\pi)$ or $[-\pi, \pi)$.[¶]

The parametrization (16) can be more useful than (14), (15) because the η_{ij} drop out of the expression $R^\dagger R$, which arises from (the often occurring) $\mathbf{Y}_\nu^\dagger \mathbf{Y}_\nu$ for degenerate \mathcal{M}_R .

A. Perturbativity

Perturbativity of the Yukawa couplings gives constraints of the form [17]

$$\text{tr}(\mathbf{Y}_\nu^\dagger \mathbf{Y}_\nu) = \frac{1}{v^2} \sum_{i=1}^n \sum_{j=1}^3 |R_{ij}|^2 M_i m_j \lesssim \mathcal{O}(1). \quad (17)$$

This inequality is satisfied if

$$|R_{ij}| \lesssim \frac{v}{\sqrt{3n M_i m_j}} \simeq 10^6 \sqrt{\frac{3}{n}} \sqrt{\frac{1 \text{ TeV}}{M_i}} \sqrt{\frac{10^{-2} \text{ eV}}{m_j}}, \quad (18)$$

which constrains each entry in R . Note that the upper bound goes down for increasing n . Once the smallest masses are specified, the above equation can be used to give a total upper bound on the parameters ρ_{ij} . For example, the largest entry in R can be estimated as $\cosh \rho_{\max}$ if just one ρ_{ij} is large, and $(\cosh \rho_{\max})^{3(n-2)}$ if all ρ_{ij} are large. This typically leads to upper bounds on $|\rho_{ij}|$ of $\mathcal{O}(1-10)$.

B. Lepton Flavor Violation

In the seesaw limit, the neutrino states ν_L that couple to the charged current are approximately given by [18] $\nu_L \simeq (\mathbb{1} - \frac{1}{2} m_D^\dagger \mathcal{M}_R^{-2} m_D) U_{\text{PMNS}} \nu_m + m_D^\dagger \mathcal{M}_R^{-1} N^c$, with the mass eigenstates ν_m and N^c . This leads to lepton flavor violating decays $\ell \rightarrow \ell' \gamma$ [19], of which $\mu \rightarrow e \gamma$ gives the strongest constraint:

$$\text{BR}(\mu \rightarrow e \gamma) \simeq \frac{3\alpha}{8\pi} \left| \left(m_D^\dagger \mathcal{M}_R^{-1} A \mathcal{M}_R^{-1} m_D \right)_{\mu e} - \frac{1}{2} \left(m_D^\dagger \mathcal{M}_R^{-2} m_D U_{\text{PMNS}} B U_{\text{PMNS}}^\dagger + \text{h.c.} \right)_{\mu e} \right|^2, \quad (19)$$

with the diagonal matrices $A_{ij} \equiv \delta_{ij} g(M_j^2/M_W^2)$ and $B_{ij} \equiv \delta_{ij} g(m_j^2/M_W^2)$ and the monotonic decreasing loop function

$$g(x) = \frac{10 - 43x + 78x^2 - 49x^3 + 4x^4 + 18x^3 \log x}{6(x-1)^4}, \quad g(0) = \frac{5}{3}, \quad g(1) = \frac{17}{12}, \quad g(x \rightarrow \infty) = \frac{2}{3}. \quad (20)$$

Here we already omitted the contribution from the light neutrinos, as the corresponding term $(U_{\text{PMNS}} B U_{\text{PMNS}}^\dagger)_{\mu e}$ is suppressed by the tiny mass-squared differences of the active neutrinos [19]. The remaining contributions are naively of order $(m_D/\mathcal{M}_R)^2$, but can be enhanced via R . In the approximation $M_j \gg M_W \gg m_k$ we have $A \propto \mathbb{1} \propto B$, so Eq. (19) simplifies to

$$\text{BR}(\mu \rightarrow e \gamma) \simeq \frac{3\alpha}{8\pi} \left| \left(m_D^\dagger \mathcal{M}_R^{-2} m_D \right)_{\mu e} \right|^2 \simeq \frac{3\alpha}{8\pi} \left| \left(U_{\text{PMNS}} \sqrt{d_m} R^\dagger \mathcal{M}_R^{-1} R \sqrt{d_m} U_{\text{PMNS}}^\dagger \right)_{\mu e} \right|^2. \quad (21)$$

[¶] The parameter space of the angles can be further reduced using discrete field redefinitions like in Refs. [16], which however lies outside the scope of this letter.




Figure 1: Distribution of the \mathcal{M}_R eigenvalues M_k according to the linear measure from Eq. (23) with the boundary constraint $\sum_{k=1}^n M_k^2 \leq M_0^2$.

Since this branching ratio can easily exceed the current limit $\text{BR}(\mu \rightarrow e\gamma) < 2.4 \times 10^{-12}$ [20]—for perturbative couplings—it gives constraints on the ρ_{ij} . Naively, the relevant product in Eq. (21) is $|R|^2 m/M$, so the LFV constraints have a different structure than the perturbativity bound (18). Note that this constraint is connected to the unitarity of the lepton mixing matrix [21], because the relevant matrix $(1 - \frac{1}{2}m_D^\dagger \mathcal{M}_R^{-2} m_D)U_{\text{PMNS}}$ is no longer unitary at this order. Consequently, the above bound also ensures the validity of our parametrization for \mathbf{Y}_ν from Eq. (12), as the matrix U_{PMNS} in \mathbf{Y}_ν only corresponds to the lepton mixing matrix if $(m_D^\dagger \mathcal{M}_R^{-2} m_D)_{ij} \ll 1$.

While not particularly nice to look at analytically, the constraints given here should be useful in numerical scans involving \mathbf{Y}_ν for arbitrary many right-handed neutrinos, extending for example the analysis of Ref. [17].

V. APPLICATION TO NEUTRINO ANARCHY

The idea that the large lepton mixing angles and small neutrino hierarchy are not due to some flavor symmetry but rather the absence of any distinction between the neutrino generations has been around for over a decade [22, 23]. The basis independence of the neutrino mass and mixing matrices leads to a distribution of the mixing angles according to the Haar measure of the Lie group G that diagonalizes \mathcal{M}_ν (so $G = U(3)$ for complex, $G = O(3)$ for real \mathcal{M}_ν). Since these distributions prefer large mixing angles, this ansatz seems to fit well to the observed large leptonic mixing angles (see however also a critique of the anarchy approach in Refs. [24, 25]). While the distribution of the mixing angles is uniquely given by the Haar measure, the distributions of the masses and Yukawa couplings are not unique, and usually the simplest (linear) measure is used (listed in App. B).

Let us now discuss neutrino anarchy, i.e. basis independence, with n right-handed neutrinos. In the seesaw limit we have the low-energy mass matrix for the active neutrinos from Eq. (11)

$$\mathcal{M}_\nu \simeq -v^2 V_L^* D_Y^T V_R^T \mathcal{M}_R^{-1} V_R D_Y V_L^\dagger, \quad (22)$$

where V_L and V_R are now assumed to be distributed according to the Haar measure of $U(3)$ and $U(n)$, respectively, while we take the simplest linear measure for the eigenvalues of \mathcal{M}_R [23]

$$d\mathcal{M}_R \propto \prod_{i < j}^n |M_j^2 - M_i^2| \prod_{k=1}^n M_k dM_k. \quad (23)$$

We employ the invariant boundary $\text{tr}(\mathcal{M}_R^\dagger \mathcal{M}_R) = \sum M_i^2 \leq M_0^2$ for the scanning region; see Fig. 1 for a visualization of the distribution.** For the singular values of D_Y we use the linear measure (see App. B

** On a technical note, random unitary matrices following the Haar distribution were obtained from a QR decomposition [26], while random entries out of the linear measures $d\mathcal{M}_R$ and dD_Y were drawn using a Metropolis algorithm [27]. This algorithm is fast even for large n and conveniently works without knowing the normalization factor of the distribution.




Figure 2: Distribution of the singular values y_k of the Yukawa matrix according to the linear measure from Eq. (24) for $n = 3$ (black) and $n = 20$ (red/dashed) with the boundary constraint $\sum_{k=1}^3 y_k^2 \leq y_0^2$.




Figure 3: $\log_{10}[R_\nu] \equiv \log_{10}[(m_2^2 - m_1^2)/(m_3^2 - m_1^2)]$ for various n . The purple/dashed vertical lines indicate the 3σ range for $R_\nu \simeq 0.03$ [28] (for NH). The black vertical line divides the NH and IH solutions.

for the derivation)

$$dD_Y \propto \prod_{m=1}^3 (y_m^2)^{n-3} \prod_{i<j}^3 (y_i^2 - y_j^2)^2 \prod_{k=1}^3 y_k dy_k, \quad (24)$$

with a similar boundary $\text{tr}(\mathbf{Y}_\nu^\dagger \mathbf{Y}_\nu) = \text{tr}(D_Y^\dagger D_Y) = \sum_j^3 y_j^2 \leq y_0^2$. The effect of large n will be a reduced hierarchy in the singular values y_i (see Fig. 2). Note that Eq. (24) is only valid for $n \geq 3$, see App. C for the used measure for $n = 2$.

The value $v^2 y_0^2 / M_0$ will fix the overall light neutrino mass scale and can be used to fix one of the mass-squared differences Δm_{ij}^2 . The PMNS mixing angles and phases do not change with n and are given by the Haar measure for $U(3)$ [23] (ignoring unphysical phases):

$$dU_{\text{PMNS}} \propto ds_{12}^2 ds_{23}^2 dc_{13}^4 d\delta d\alpha d\beta, \quad (25)$$

so the only effect of $n \neq 3$ is a change of the three eigenvalues of \mathcal{M}_ν , i.e. the ratio $R_\nu \equiv (m_2^2 - m_1^2)/(m_3^2 - m_1^2)$, where we sorted the masses like $m_1 \leq m_2 \leq m_3$. In this notation, $R_\nu < 1/2$ corresponds to normal hierarchy (NH) and $R_\nu > 1/2$ to inverted hierarchy (IH), with best-fit values $R_\nu^{\text{NH}} = 0.03$ and $R_\nu^{\text{IH}} = 0.97$ (IH), respectively, taken from a recent global fit [28].

We observe from Fig. 3 that for increasing n , the distribution for R_ν shifts to larger values, while the width decreases. This reduction of hierarchy can be tracked back mainly to the Yukawa couplings, as the factor $\prod_{m=1}^3 (y_m^2)^{n-3}$ in dD_Y pulls the y_i tightly together. An analysis without this factor shows that the behavior of R_ν goes in the opposite direction, i.e. the maximum shifts towards small values for large n while being diluted. The distribution then however quickly converges as the only n dependent change

comes from $V_R^T \mathcal{M}_R^{-1} V_R$, for which only the first dozen or so M_i are relevant, due to the hierarchy in \mathcal{M}_R . Correspondingly, to shift the maximum of R_ν to its observed value, one not only has to omit the factor $\prod_{m=1}^3 (y_m^2)^{n-3}$ in dD_Y , but also increase the hierarchy by inserting something like $\prod_{i < j} (y_i^2 - y_j^2)^2$, i.e. using a non-linear measure (see Ref. [6] for other y_i distributions that accomplish this task). We conclude that the large n limit in the anarchy approach worsens the agreement with data. While the one data point Nature provides can obviously not be used to find the overlying distribution, it should be fair to say that anarchy works best for small n .

Let us finally comment on the limit $n \rightarrow \infty$. As the Yukawa couplings y_i become quasi-degenerate for very large n , the neutrino masses are simply given by the upper left 3×3 submatrix of $V_R^T \mathcal{M}_R^{-1} V_R$ (times a prefactor). Since there is no preferred direction in the V_R rotations, the entries $(V_R)_{ij}$ have the same mean magnitude for a given n . The mean entries $\langle (V_R^T \mathcal{M}_R^{-1} V_R)_{jk} \rangle$ then take the form of a random walk with decreasing step-size $\sum_m^n e^{i\theta_m}/M_m$, and due to the hierarchy in the singlet masses M_k , only the first couple of steps are relevant. The mean magnitudes are again the same for all entries, so the only structure in the neutrino mass matrix comes from random phases $\tilde{\theta}_{jk}$: $\langle (V_R^T \mathcal{M}_R^{-1} V_R)_{jk} \rangle \propto e^{i\tilde{\theta}_{jk}}$. It is easy to show that a matrix of this type predominantly yields NH solutions, so even though IH becomes more and more probable for large n , it will never dominate.

VI. CONCLUSION

One of the simplest ways to explain the masses of the active neutrinos is the introduction of right-handed partners. As the number n of these SM singlets is in principle unconstrained, it is interesting to study implications of varying n . We have given explicit parametrizations for the mixing and coupling matrices connecting the n right-handed neutrinos to the SM in terms of physical parameters, both in the top-down and the bottom-up approach (Casas-Ibarra-parametrization). For the latter, constraints on the involved parameters from perturbativity of the Yukawa couplings as well as charged lepton flavor violation have been discussed.

As a novel application of the n singlet framework we studied basis independence in the neutrino sector, i.e. anarchy. Of the low-energy neutrino parameters, only the neutrino mass distribution changes with n , and we showed that anarchy—with the simplest linear measure—seems to prefer small n in view of the observed mass-squared differences.

Acknowledgments

The author thanks Werner Rodejohann and Mattias Blennow for reading the manuscript and valuable comments. This work was supported by the the ERC under the Starting Grant MANITOP and by the IMPRS for Precision Tests of Fundamental Symmetries.

Appendix A: The Group $U(N)$

For convenience we provide a short and rather colloquial review of the group $U(N, \mathbb{C}) \equiv U(N)$. Formally one can define it as the set of complex $N \times N$ matrices that leave the inner product of \mathbb{C}^N invariant, i.e. $U(N) = \{V \in \mathbb{C}^{N \times N} \mid V^\dagger V = \mathbb{1}\}$. With a little effort one can further show that the entries of $U(N)$ form a group under matrix multiplication, actually even a compact connected Lie group. Counting the number of entries in $V \in U(N)$ and constraints, we find that $U(N)$ has N^2 real parameters. One can show that every element in $U(N)$ can be written as $V = \exp(A)$ with skew-hermitian $A = -A^\dagger$. We can choose a basis for $A = \sum \omega_j X_j$ such that the N^2 linearly independent X_j satisfy Lie algebra relations $[X_k, X_l] = if_{klm} X_m$ with structure constants f_{klm} .

The N^2 generators X_j can be further separated into $N(N+1)/2$ symmetric generators $X_s = iY = iY^T$ and $N(N-1)/2$ antisymmetric real generators $X_a = -X_a^T$. It also proves convenient to treat the N diagonal (symmetric) generators Y_d separately, as they are especially easy to exponentiate. To fix the normalization, we chose $(Y_d^k)_{ij} = \delta_{ij} \delta_{jk}$, while the $N(N-1)/2$ non-diagonal Y_s and X_a have just two non-vanishing entries, namely $+1$ in the upper right half and ± 1 in the lower left (plus sign for Y_s , minus sign for X_a). Note that this normalization—and therefore also f_{klm} —differs from the common

convention, but is of no importance in the following (see also Refs. [9]). A general element of $U(N)$ can then be written as the product of all rotations:

$$U(N) \ni V = \prod_{j=1}^N \exp(i\alpha_j Y_d^j) \prod_{k=1}^{N(N-1)/2} \exp(i\beta_k Y_s^k) \prod_{m=1}^{N(N-1)/2} \exp(\gamma_m X_a^m). \quad (\text{A1})$$

The equivalence of this parametrization to the initial $V = \exp(A)$ can be proven using the Baker-Campbell-Hausdorff formula and the fact that the generators satisfy a Lie algebra. The order of the rotations in Eq. (A1) is of course arbitrary [29] and should be chosen to simplify given expressions. The three different types of rotations take the form

$$e^{i\beta Y_s} = \begin{pmatrix} 1 & & & & \\ & \ddots & & & \\ & & \cos(\beta) & i \sin(\beta) & \\ & & i \sin(\beta) & \cos(\beta) & \\ & & & & \ddots & \\ & & & & & 1 \end{pmatrix}, \quad e^{\gamma X_a} = \begin{pmatrix} 1 & & & & \\ & \ddots & & & \\ & & \cos(\gamma) & \sin(\gamma) & \\ & & -\sin(\gamma) & \cos(\gamma) & \\ & & & & \ddots & \\ & & & & & 1 \end{pmatrix}, \quad (\text{A2})$$

and

$$\exp(i\alpha Y_d) = \text{diag}(1, \dots, 1, \exp(i\alpha), 1, \dots, 1), \quad (\text{A3})$$

with real angles α, β, γ . Another useful parametrization for a general unitary matrix along the same lines as above is given by

$$U(N) \ni V = \prod_{j=1}^N \exp(i\alpha'_j Y_d^j) \prod_{k=1}^{N(N-1)/2} \exp(i\beta'_k Y_s^k + \gamma'_k X_a^k), \quad (\text{A4})$$

where we combined the generators X_a and Y_s that have the same vanishing entries. This yields a product of matrices of the type

$$\exp(i\beta'_j Y_s^j + \gamma'_j X_a^j) = \begin{pmatrix} 1 & & & & \\ & \ddots & & & \\ & & \cos(z_j) & \sin(z_j)e^{i\xi_j} & \\ & & -\sin(z_j)e^{-i\xi_j} & \cos(z_j) & \\ & & & & \ddots & \\ & & & & & 1 \end{pmatrix}, \quad (\text{A5})$$

with the real parameters $z_j = |u_j|$, $\xi_j = \arg(u_j)$ and $u_j = i\beta'_j - \gamma'_j$ in our chosen normalization. Once again, the ordering of the matrices in Eq. (A4) is arbitrary.

Appendix B: Measures for Anarchy

This appendix provides the necessary measures for the anarchy approach in Sec. V. We refer to Ref. [23] for a detailed derivation of the known measures, and simply quote the result for real and complex Majorana $N \times N$ matrices:

$$d\mathcal{M}_{\text{real}} \propto \prod_{i < j}^N |D_i - D_j| \prod_{k=1}^N dD_k dO, \quad d\mathcal{M}_{\text{complex}} \propto \prod_{i < j}^N |D_i^2 - D_j^2| \prod_{k=1}^N D_k dD_k dU, \quad (\text{B1})$$

where the decompositions $\mathcal{M}_{\text{real}} = ODO^T$ and $\mathcal{M}_{\text{complex}} = UDU^T$ were used. dO and dU denote the usual Haar measure for $O(N)$ and $U(N)$, respectively, while the diagonal matrix D contains the eigenvalues of \mathcal{M} . We will omit the calculation of the normalization factor for all measures in this appendix, as they are irrelevant for our considerations. It can in principle be calculated by integration over some volume, e.g. $\text{tr}(\mathcal{M}_{\text{complex}}^\dagger \mathcal{M}_{\text{complex}}) \leq M_0^2$, up to the arbitrary overall mass scale M_0 .

For Dirac matrices, the measures read [23]

$$dm_{\text{real}} \propto \prod_{i<j}^N |D_i^2 - D_j^2| \prod_{k=1}^N dD_k dO_L dO_R, \quad dm_{\text{complex}} \propto \prod_{i<j}^N (D_i^2 - D_j^2)^2 \prod_{k=1}^N D_k dD_k dU_L dU_R, \quad (\text{B2})$$

with $m_{\text{real}} = O_R D O_L^T$, $m_{\text{complex}} = U_R D U_L^\dagger$, and the Haar measures for $O(N)$ $dO_{L,R}$ and $U(N)$ $dU_{L,R}$ respectively. For dm_{complex} , unphysical phases in the overlap of dU_L and dU_R should be modded out, once a parametrization is specified.

These invariant measures are of course not unique, but rather the simplest consistent ansatz that give the right mass dimension.^{††} With the above equations, one can easily write down the measure for an hermitian matrix $H = U h U^\dagger = H^\dagger$ under $U(N)$:

$$dH \propto \prod_{i<j}^N (h_i - h_j)^2 \prod_{k=1}^N dh_k dU. \quad (\text{B3})$$

As a cross check, one can plug the hermitian matrix $m_{\text{complex}}^\dagger m_{\text{complex}}$ into Eq. (B3) to rederive Eq. (B2). We can now give a measure for a non-quadratic $n \times 3$ complex Dirac matrix $M_{SVD} = U_R D_Y U_L^\dagger$ with $U_R \in U(n)$, $U_L \in U(3)$. Considering $M_{SVD}^\dagger M_{SVD}$ shows that dM_{SVD} should at least contain a factor

$$\prod_{i<j}^3 (y_i^2 - y_j^2)^2 \prod_{k=1}^3 y_k dy_k, \quad (\text{B4})$$

which however has too small of a mass dimension. The same procedure for the $n \times n$ hermitian $M_{SVD} M_{SVD}^\dagger$ runs into problems, because $n - 3$ of the eigenvalues are zero and would give a vanishing measure. Taking the product only over the non-vanishing factors however yields

$$dM_{SVD} \propto \prod_{m=1}^3 (y_m^2)^{n-3} \prod_{i<j}^3 (y_i^2 - y_j^2)^2 \prod_{k=1}^3 y_k dy_k dU_L dU_R, \quad (\text{B5})$$

which has the right mass dimension ($6n$) and can therefore be viewed as the linear measure for M_{SVD} . The overlap of dU_L and dU_R —this time including angles, see Sec. III—should again be modded out in a given parametrization to avoid overcounting of physically equivalent configurations (similar to gauge-fixing, as pointed out in Ref. [25]).

The measure (B5) can of course be easily extended to a general measure over $O(N) \times O(M)$ or $U(N) \times U(M)$ with $M \leq N$:

$$dM_{\text{real}} \propto \prod_{i<j}^M |y_i^2 - y_j^2| \prod_{k=1}^M y_k^{N-M} dy_k dO_L dO_R, \quad dM_{\text{complex}} \propto \prod_{i<j}^M (y_i^2 - y_j^2)^2 \prod_{k=1}^M y_k^{1+2(N-M)} dy_k dU_L dU_R, \quad (\text{B6})$$

with the $N \times M$ matrices $M_{\text{real}} = O_R D_Y O_L^T$ and $M_{\text{complex}} = U_R D_Y U_L^\dagger$. Unphysical rotations need to be modded out again. A simple check shows that the mass dimensions are right and $N = M$ indeed leads back to Eq. (B2). Eq. (B6) can actually be obtained from exactly these requirements, without

^{††} For example, $\mathcal{M}_{\text{complex}}$ contains $N(N+1)$ independent real parameters, so the measure $d\mathcal{M}_{ij}$ has mass dimension $N(N+1)$. Rewriting $d\mathcal{M}_{ij}$ in the form (B1) matches this mass dimension, as can be easily verified.

the need to use hermitian matrices in the derivation. For example, dM_{complex} should contain the known $N = M$ factor $\prod_{i < j}^M (y_i^2 - y_j^2)^2 \prod_{k=1}^M dy_k$, which leaves some factor of mass dimension $2M$ to the power of $N - M$. Since a factor $\prod_{i < j}^M (y_i^k - y_j^k)^m$ —which would increase the hierarchy in the y_i —has mass dimension $M(M-1)km/2$, it can only be used in special cases, not for general M (at least for integer k and m). Thus, for general M , the only factor with mass dimension $2M$ is $\prod_i^M y_i^2$, leading back to Eq. (B6).

Appendix C: The Case $n = 2$

In this appendix we collect the formulae for the case $n = 2$ (see also Refs. [3, 30]), which is a little different than the $n \geq 3$ cases and would interrupt the flow if included in the main text. With the 2×2 diagonal matrix \mathcal{M}_R we find the top-down parametrization in the singular value decomposition for the Yukawa coupling

$$\mathbf{Y}_\nu = V_R \begin{pmatrix} y_1 & 0 & 0 \\ 0 & y_2 & 0 \end{pmatrix} V_L^\dagger. \quad (\text{C1})$$

Two phases of V_R can be absorbed by V_L^\dagger , three of V_L in the lepton fields. Since one of the diagonal phase rotations of V_L acts on the zeros, there are only two phases in V_L left, so we can write

$$V_R = \begin{pmatrix} \cos(\alpha_{12}) & \sin(\alpha_{12})e^{i\xi_{12}} \\ -\sin(\alpha_{12})e^{-i\xi_{12}} & \cos(\alpha_{12}) \end{pmatrix}, \quad (\text{C2})$$

$$V_L = \begin{pmatrix} c_{12}^L c_{13}^L & s_{12}^L c_{13}^L & s_{13}^L e^{-i\beta_1^L} \\ -c_{23}^L s_{12}^L - s_{23}^L s_{13}^L c_{12}^L e^{i\beta_1^L} & c_{23}^L c_{12}^L - s_{23}^L s_{13}^L s_{12}^L e^{i\beta_1^L} & s_{23}^L c_{13}^L \\ s_{23}^L s_{12}^L - c_{23}^L s_{13}^L c_{12}^L e^{i\beta_1^L} & -s_{23}^L c_{12}^L - c_{23}^L s_{13}^L s_{12}^L e^{i\beta_1^L} & c_{23}^L c_{13}^L \end{pmatrix} \text{diag}(1, e^{i\beta_2^L/2}, 1), \quad (\text{C3})$$

with the shorthand notation $s_{ij}^L = \sin \theta_{ij}^L$ etc. for the mixing angles. The linear measure for $y_{1,2}$ for the anarchy framework can be derived along the same lines as in App. B, with the result

$$dD_Y \propto (y_1^2 - y_2^2)^2 y_1^3 y_2^3 dy_1 dy_2. \quad (\text{C4})$$

Finally, the bottom-up parametrization is given by Eq. (12) with two different R matrices depending on the low-energy neutrino hierarchy ($\gamma \in \mathbb{C}$) [31]:

$$R_{\text{NH}} = \begin{pmatrix} 0 & \cos \gamma & -\sin \gamma \\ 0 & \sin \gamma & \cos \gamma \end{pmatrix}, \quad R_{\text{IH}} = \begin{pmatrix} \cos \gamma & -\sin \gamma & 0 \\ \sin \gamma & \cos \gamma & 0 \end{pmatrix}. \quad (\text{C5})$$

-
- [1] P. Minkowski, Phys. Lett. B **67**, 421 (1977);
M. Gell-Mann, P. Ramond, and R. Slansky, in *Supergravity*, p. 315, edited by P. van Nieuwenhuizen and D. Freedman, North Holland, Amsterdam, 1979;
T. Yanagida, Proc. of the *Workshop on Unified Theory and the Baryon Number of the Universe*, KEK, Japan, 1979;
R. N. Mohapatra and G. Senjanović, Phys. Rev. Lett. **44**, 912 (1980).
 - [2] M. Fukugita and T. Yanagida, Phys. Lett. B **174**, 45 (1986).
 - [3] See, for example, S. F. King, Nucl. Phys. B **576**, 85 (2000) [arXiv:hep-ph/9912492];
P. H. Frampton, S. L. Glashow and T. Yanagida, Phys. Lett. B **548**, 119 (2002) [arXiv:hep-ph/0208157];
S. Antusch, P. Di Bari, D. A. Jones and S. F. King, arXiv:1107.6002 [hep-ph];
K. Harigaya, M. Ibe and T. T. Yanagida, arXiv:1205.2198 [hep-ph].
 - [4] M. T. Eisele, Phys. Rev. D **77**, 043510 (2008) [arXiv:0706.0200 [hep-ph]].
 - [5] J. R. Ellis and O. Lebedev, Phys. Lett. B **653**, 411 (2007) [arXiv:0707.3419 [hep-ph]].
 - [6] B. Feldstein and W. Klemm, Phys. Rev. D **85**, 053007 (2012) [arXiv:1111.6690 [hep-ph]].
 - [7] W. Buchmüller, K. Hamaguchi, O. Lebedev, S. Ramos-Sánchez and M. Ratz, Phys. Rev. Lett. **99**, 021601 (2007) [arXiv:hep-ph/0703078].

- [8] K. R. Dienes, E. Dudas and T. Gherghetta, Nucl. Phys. B **557**, 25 (1999) [arXiv:hep-ph/9811428];
G. R. Dvali and A. Y. Smirnov, Nucl. Phys. B **563**, 63 (1999) [arXiv:hep-ph/9904211];
N. Arkani-Hamed, S. Dimopoulos, G. R. Dvali and J. March-Russell, Phys. Rev. D **65**, 024032 (2002) [arXiv:hep-ph/9811448].
- [9] J. Schechter and J. W. F. Valle, Phys. Rev. D **22**, 2227 (1980);
G. C. Branco, L. Lavoura and M. N. Rebelo, Phys. Lett. B **180**, 264 (1986).
- [10] For recent reviews, see W. Rodejohann, Int. J. Mod. Phys. E **20**, 1833 (2011) [arXiv:1106.1334 [hep-ph]];
J. J. Gómez-Cadenas, J. Martín-Albo, M. Mezzetto, F. Monrabal and M. Sorel, Riv. Nuovo Cim. **35**, 29 (2012) [arXiv:1109.5515 [hep-ex]].
- [11] A. Broncano, M. B. Gavela and E. E. Jenkins, Phys. Lett. B **552**, 177 (2003) [Erratum-ibid. B **636**, 330 (2006)] [arXiv:hep-ph/0210271].
- [12] J. A. Casas and A. Ibarra, Nucl. Phys. B **618**, 171 (2001) [arXiv:hep-ph/0103065];
J. A. Casas, A. Ibarra and F. Jiménez-Alburquerque, JHEP **0704**, 064 (2007) [arXiv:hep-ph/0612289].
- [13] A. de Gouvêa and W. C. Huang, Phys. Rev. D **85**, 053006 (2012) [arXiv:1110.6122 [hep-ph]];
P. S. B. Dev, R. Franceschini and R. N. Mohapatra, arXiv:1207.2756 [hep-ph].
- [14] R. N. Mohapatra, Phys. Rev. Lett. **56**, 561 (1986);
R. N. Mohapatra and J. W. F. Valle, Phys. Rev. D **34**, 1642 (1986).
- [15] E. K. Akhmedov, M. Lindner, E. Schnapka and J. W. F. Valle, Phys. Rev. D **53**, 2752 (1996) [arXiv:hep-ph/9509255];
S. M. Barr, Phys. Rev. Lett. **92**, 101601 (2004) [arXiv:hep-ph/0309152];
C. H. Albright and S. M. Barr, Phys. Rev. D **69**, 073010 (2004) [arXiv:hep-ph/0312224].
- [16] A. de Gouvêa and J. Jenkins, Phys. Rev. D **78**, 053003 (2008) [arXiv:0804.3627 [hep-ph]];
M. Blennow and E. Fernández-Martínez, Phys. Lett. B **704**, 223 (2011) [arXiv:1107.3992 [hep-ph]].
- [17] J. A. Casas, J. M. Moreno, N. Rius, R. Ruiz de Austri and B. Zaldivar, JHEP **1103**, 034 (2011) [arXiv:1010.5751 [hep-ph]].
- [18] W. Grimus and L. Lavoura, JHEP **0011**, 042 (2000) [arXiv:hep-ph/0008179].
- [19] T. P. Cheng and L. F. Li, Phys. Rev. Lett. **45**, 1908 (1980).
- [20] J. Adam *et al.* [MEG Collaboration], Phys. Rev. Lett. **107**, 171801 (2011) [arXiv:1107.5547 [hep-ex]].
- [21] Thorough discussions can be found in S. Antusch, C. Biggio, E. Fernández-Martínez, M. B. Gavela and J. López-Pavón, JHEP **0610**, 084 (2006) [arXiv:hep-ph/0607020];
D. V. Forero, S. Morisi, M. Tórtola and J. W. F. Valle, JHEP **1109**, 142 (2011) [arXiv:1107.6009 [hep-ph]].
- [22] L. J. Hall, H. Murayama and N. Weiner, Phys. Rev. Lett. **84**, 2572 (2000) [arXiv:hep-ph/9911341];
G. Altarelli, F. Feruglio and I. Masina, JHEP **0301**, 035 (2003) [arXiv:hep-ph/0210342];
A. de Gouvêa and H. Murayama, Phys. Lett. B **573**, 94 (2003) [arXiv:hep-ph/0301050];
A. de Gouvêa and H. Murayama, arXiv:1204.1249 [hep-ph].
- [23] N. Haba and H. Murayama, Phys. Rev. D **63**, 053010 (2001) [arXiv:hep-ph/0009174].
- [24] M. Hirsch, arXiv:hep-ph/0102102.
- [25] J. R. Espinosa, arXiv:hep-ph/0306019.
- [26] A. Edelman and N. R. Rao, Acta Numerica **14**, 233-297 (2005);
F. Mezzadri, Notices of the AMS, Vol. **54**, Issue 5, 592-604 (2007) [arXiv:math-ph/0609050].
- [27] N. Metropolis, A. W. Rosenbluth, M. N. Rosenbluth, A. H. Teller and E. Teller, J. Chem. Phys. **21**, 1087 (1953).
- [28] D. V. Forero, M. Tórtola and J. W. F. Valle, arXiv:1205.4018 [hep-ph].
- [29] M. Gronau, R. Johnson and J. Schechter, Phys. Rev. D **32**, 3062 (1985).
- [30] Early references include A. Y. Smirnov, Phys. Rev. D **48**, 3264 (1993) [arXiv:hep-ph/9304205];
A. Kleppe, in *Lohusalu 1995, Neutrino physics* pp. 118–125;
E. Ma, D. P. Roy and U. Sarkar, Phys. Lett. B **444**, 391 (1998) [arXiv:hep-ph/9810309].
- [31] A. Ibarra and G. G. Ross, Phys. Lett. B **591**, 285 (2004) [arXiv:hep-ph/0312138].